
Argonne HEP Division DOE Review : May 12, 2004

ARGONNE HEP THEORY GROUP
OVERVIEW

C.E.M. Wagner

HEP Division

Argonne National Laboratory

- Theory at Argonne
- External Funding Awards
- Some research Highlights
- HEP Community Activities
- Outlook

Composition of the Group

- The HEP Theory Group has Five Permanent Staff Members:
Ed Berger, Geoff Bodwin, Don Sinclair,
Carlos Wagner, Cosmas Zachos.
- It has, at this point, **no** Junior Scientist Member.
- The activities of the Group have been reinforced by a number of young postdoctoral appointees.
- The number of regular postdoctoral fellows supported by the HEP Division in the last years has fluctuated around three (going down to two next year).
- Additional postdoctoral appointments, however, have been achieved by the acquisition of Grants, or by agreements between the Laboratory and the Univ. of Chicago.

University Affiliations and Teaching Activities

Members of the Group are affiliated with Universities in the Chicagoland area:

Ed Berger – Adjunct Professor, Michigan State University, since 1997.

Carlos Wagner – Associate Professor, Univ of Chicago. Joint appointment with Argonne National Laboratory, since 1999.

Teaching Advanced Classical Electrodynamics during the Spring Quarter 2003.

Introduction to Cosmology, Spring Quarter 2004.

Close relationship with the University of Chicago was used to create a collaborative effort between ANL and the Univ. of Chicago, that led to the appointments of D. Kaplan, C. Chiang, G. Servant, and **Irina Mocioiu**.

Theory at Argonne

- Effort is being made to increase the visibility of the group through the organization of workshops and meetings.
- Since May, 2003, members of the group have organized:
Workshop on QCD on Extreme Environments, June 29–July 3, 2004, <http://www.hep.anl.gov/dks/qcdxe/>
Workshop on Supersymmetry, Extra Dimensions, and Higgs Bosons, May 24–28, 2004 ,
<http://gate.hep.anl.gov/berger/ANLWorkshop2004/>
Workshop on Branes and Generalized Dynamics, October 20–24, 2003,
<http://www.hep.anl.gov/czachos/ANLworkshop.html>
Workshop on Trends in Neutrino Physics, May 12–16, 2003 , <http://www.neutrinooscillation.org/trends.html>
- There is also an effort to increase interactions among different theory groups at ANL : **Ed Berger** organized a February 16, 2004 **Argonne Lab-wide Theory Afternoon**.
- **Ed Berger** is the Co-Director of the LDRD funded Argonne Lab-wide Theory Institute.
- **C. Wagner** has been acting as the head of the **ANL Theory Committee**.

Recent ANL and External Funding Awards

- **Argonne Theory Institute 2004**
Supersymmetry, Extra Dimensions, and Higgs Bosons
\$20K awarded to E. Berger and C. Wagner, to organize a workshop to take place May 24–28, 2004.
- **Argonne Theory Institute 2004**
QCD in Extreme Environments
\$10K awarded to D. Sinclair, to organize a workshop to take place on June 29–July 3, 2004.
- **Argonne Theory Institute 2003**
Trends in Neutrino Physics: \$20K awarded for a workshop May 12–16, 2003.
- **Laboratory Graduate Student Program: Support for D. Morrissey, to do his thesis under the supervision of C. Wagner at the Theory Group of the HEP Division, \$24K awarded for the first year, starting in April 2003.**
- **Argonne Individual Investigator Award, October 2001,**
“Extra Space-Time Dimensions”, \$ 57 K awarded to E. Berger and C. Wagner. Supported the appointment of B. Murakami for two years.

Response to DOE Recommendations

Joint Junior Faculty Position with the Univ. of Chicago

- Idea: Take advantage of new program to create joint junior faculty position in HEP phenomenology.
- Initiative presented to the Univ. of Chicago Physics Department. Voted favorably in an unanimous way (Vote: 20 to nothing).
- Search Committee: Ed Berger, Carlos Wagner, Jon Rosner and Emil Martinec.
- Initiative abandoned last year, due to a senior faculty search at the University of Chicago.
- Revived this year with an excellent candidate, but UofC physics department did not support the idea with the necessary strength to make it work.
- HEP Division still working very hard to get a junior faculty in Theory, either jointly with local universities (UofC, Northwestern), or by genuine Division funds.

Career Paths

Among the postdocs who worked at Argonne in the last ten years, many have been hired by Laboratories:

J. Hewett (1993), Faculty at SLAC; S. Mrenna (1999), Junior Faculty at Fermilab; Z. Sullivan (2001), T.M.P. Tait (2002), Postdocs at Fermilab; J. Campbell, Fellow at CERN (2004)

Other recent postdocs obtained Positions at Universities

I. Knowles (1993), Glasgow U; R. Kauffman (1993) Muhlenberg College; C. Coriano (1995), Lecce U; S. Kim (1995) Sejong U; M. Klasen (1999) Grenoble U.; B. Harris (2001), Robert Morris U; D.E. Kaplan (2001), John Hopkins U; G. Servant (2003), Saclay; C.W. Chiang (2003), Nat. Central U, Taiwan; J. Lee (2003) Korea University; J. Jiang, Postdoc at Oregon U. (2004)

Some of them, explore other career paths, including

L. Gordon (1997), M. Wusthoff (1997); H. Contopanagos (1997), J. Lagae (1999), A. Petrelli (1999), G. Chalmers (2001).

High Energy Physics Community Activities in the past year

Members of the Group have taken part in numerous organizing activities and have been members of various HEP groups:

Edmond L. Berger

- Organizer, Aspen Winter Conference on Particle Physics, “Where We Are and Where We Are Going”, Aspen Center for Physics, Aspen, CO, February 1–7, 2004.
(<http://gate.hep.anl.gov/berger/Aspen04>)
- Co-Organizer, 2004 Theory Institute on Supersymmetry, Extra-Dimensions, and Higgs Boson Physics, Argonne National Laboratory, May 24–28, 2004.
(<http://gate.hep.anl.gov/berger/ANLWorkshop2004>)
- Organizer, Argonne Lab-wide Theory Afternoon, Argonne National Laboratory, February 16, 2004.
- Member, Committee on International Scientific Affairs (CISA), American Physical Society, 2003–.
- Member, Andrew Gemant Award Committee, American Institute of Physics, 2002–.
- Member, American Linear Collider Working Group, 2002–.
- Member, Coordinated Theoretical Experimental Project on QCD (CTEQ) Collaboration
- Scientific Advisory Board, Argonne National Laboratory Theory Institute, 2003–.

- Scientific Program Organizing Committee, XXXIXth Rencontres de Moriond, QCD and High Energy Hadronic Interactions, La Thuile, March 28–April 4th 2004.
- Organizing Committee and Convener of the session on the Polarized Gluon Density, Fourth Circum-Pan-Pacific Symposium on High Energy Spin Physics Univ. of Washington, Seattle, August 4–7, 2003.
- Organizing Committee, 8th Conference on the Intersections of Particle and Nuclear Physics (CIPANP 2003), New York, May 19 - 24, 2003.
- Co-organizer, Argonne Theory Institute on Trends in Neutrino Physics, May 12 - 16, 2003.
- Leader, North American Node, Quarkonium Working Group, QWNET 2003.
- Scientific Program Committee, Vth Rencontres du Vietnam, Hanoi, Vietnam, August 6–11, 2004.
- Organizing Committee, 9th Conference on the Intersections of Particle and Nuclear Physics (CIPANP 2005), 2005
- Steering Committee, International Conference on Flavor Physics 2005, (ICFP 2005), Taiwan.

Geoff Bodwin:

- Member, Local Organizing Committee, LATTICE 2004, the XXII International Symposium on Lattice Field Theory, Fermilab, June 21–26, 2004.
- Convener, Production Section, Quarkonium Working Group QWGNET Proposal to fund a Marie Curie Research Training Network on Heavy Quarkonium 2003–present.
- Member, Local Organizing Committee, Second International Workshop on Heavy Quarkonium, Fermilab, September 20-22, 2003.
- Member, Quarkonium Working Group 2002–present.
- Member, Working Group on Heavy Flavors, Workshops on Hard Probes in Heavy Ion Collisions, CERN, 2001–2003.

Don Sinclair :

- Member of NERSC users group executive committee (NUGEX).
- Member of the Quarkonium Working Group.
- Member of the Lattice SciDAC project.
- Member of the local organizing committee for Lattice2004, Fermilab.
- Organizer of Workshop on QCD in Extreme Environments.

Carlos E.M. Wagner :

- Co-organizer, with J. Terning and D. Zeppenfeld, TASI lectures 2004, “Physics in $D \geq 4$ ”, Boulder, CO, June 2004
- Co-organizer, with Wilfried Buchmuller and Gordy Kane, Workshop on Baryogenesis, Univ. of Michigan, Ann Arbor, June 2003.
- Co-organizer, with Ed Berger and Maury Goodman, Argonne Theory Institute on Neutrino Physics, May 12 - 16, 2003.
- Co-organizer, with Cosmas Zachos and Tom Cultright, Argonne Theory Workshop on Branes and Generalized Dynamics, October 20-24, 2003.
- Organizing Committee, Theory Institute on Supersymmetry, Higgs and Extra Dimensions, Argonne, May 24–28, 2004.
- Member, American Linear Collider Physics Group, 2002–
- Member, LEP Higgs Working Group, 1996–
- Head, Argonne Theory Committee, April 2003—
- Head, Theory Group, HEP Division, Sep. 2002–

Cosmas Zachos

- Principal Organizer of the Argonne HEP “Branes and Generalized Dynamics Workshop”, October 20-24, 2003, and Editor of its E-proceedings, <http://www.hep.anl.gov/czachos/ANLworkshop.html>
- Member of the Advisory Panel of J Phys A: Math Gen (IOP).
- Member of the Organizing Committee for the 2004 Coral Gables Conference, [CGC 2004], 15 - 19 December, Key Biscayne, Florida “Celebrating 40 Years of Quarks and Coral Gables Conferences”
- External Ph D Thesis Examiner for J G Wood, Univ of Queensland, Australia


Research Highlights

Transverse Momentum Distribution for Υ Production

E. L. Berger, J. W. Qiu, and Y. Wang

[hep-ph/0404158], submitted to Phys Rev D

- Fixed order QCD perturbation theory for $p\bar{p} \rightarrow \Upsilon X$ is divergent at small p_T ; $\sigma \propto (1/p_T^2) \ln^2(m_\Upsilon^2/p_T^2)$.
- Develop and justify the validity of all-orders in α_s resummation of large logs from multiple soft gluon radiation to compute the distribution at small $p_T < M_\Upsilon$.
- Demonstrate that the p_T distribution at small p_T is dominated by the region of small impact parameter, and that the distribution may be computed reliably at small p_T in resummed perturbation theory.




- Excellent agreement with 2002 CDF data over the full range of p_T .

Higgs Boson Plus 2 Jet Production: WBF Signal at NLO and QCD Backgrounds

E. L. Berger and J. Campbell

[hep-ph/0403194], submitted to Phys Rev D

- **Motivation:** after discovery of the Higgs boson, the goal becomes the measurement of its couplings. How well can this job be done in the weak boson fusion (WBF) sample?
- Independent QCD calculation of $H + 2$ jet processes
 - to gauge the effectiveness of cuts used to select the WBF signal, and
 - to evaluate the accuracy with which couplings g_H can be determined in experiments at the CERN LHC.
- As a function of the rapidity of a tagging jet, the figures show the **WBF signal** and the QCD $H + 2$ jet backgrounds, along with estimated uncertainties.



- Estimate $\delta g_H / g_H \sim 10\%$ should be possible after 200 fb^{-1} . Les Houches 2003 estimates of expected uncertainties are too optimistic ($\sim \times 2$).

Associated Production of a Top Quark and a Charged Higgs Boson tH^+


E. L. Berger, T. Han, J. Jiang, and T. Plehn

[hep-ph/0312286], submitted to Phys Rev D


- **Motivation:** identification of a charged Higgs boson H^\pm would provide evidence for a Higgs sector beyond the SM, meaning at least two Higgs doublets and, possibly, a supersymmetric Higgs sector.
- Compute the NLO inclusive and differential cross sections for $pp \rightarrow tH^- X$ at Tevatron and LHC energies in QCD and SUSY-QCD.
- For $m_H < m_{\text{top}}$, include $t\bar{t}$ pair production, with $t \rightarrow bH^+$.
- All results are fully differential, permitting selections on the momenta of both the top quark and the charged Higgs boson, plus the **study of correlations** between final state particles. The two-cutoff phase-space slicing method is used to deal with the usual soft and collinear singularities.

tH^+ Associated Production, continued

- Total cross section at the LHC vs m_H . NLO result is the dashed upper curve. Also shown is the cross section for $pp \rightarrow t\bar{t}^*; \bar{t}^* \rightarrow \bar{b}H^-$ (at small m_H).



- Correlation between $\Sigma p_T = (\vec{p}_t + \vec{p}_H)_T$ and $\Delta\phi = |\phi_t - \phi_H|$; $m_H = 250 \text{ GeV}$.




Lower limits on R-parity-violating couplings in SUSY models with light squarks

E. L. Berger and Z. Sullivan

[hep-ph/0310001], Phys Rev Letters, in press

- If R parity (R_p) is conserved, SUSY particles are produced in pairs, each of which decays to a final state that includes a stable lightest SUSY particle (LSP).
- If R_p is not conserved, SUSY particles may decay into SM particles; missing energy signatures may be lost; and SUSY may not provide a dark matter candidate.
- Interpreted the results of (negative) LEP searches for stable strongly interacting massive particles to place absolute lower limits on R_p -violating couplings:

λ'' or $\lambda' > 10^{-8} - 10^{-7}$, if $m_{\tilde{q}} < 100$ GeV.




- If R_p violation is this large, $\tilde{\chi}^0$ cannot explain dark matter.

Squark Mixing in e^+e^- Reactions

E. L. Berger, J. Lee, and T. Tait

[hep-ph/0306110], Phys Rev D **69**, 055003 (2004)


- For each squark flavor, there are two mass eigenstates, one light \tilde{q}_1 and one heavy \tilde{q}_2 . The **mixing** that results in these two states is characterized by a **mixing angle** θ_q .
- Examine prospects for **direct** measurement of the top-squark and bottom-squark mixing angles in associated production $e^+e^- \rightarrow \tilde{q}_1\tilde{q}_2^*$.
- Compute cross sections for LEP-II and a Linear Collider.
- In the context of a **light bottom squark scenario**, show that existing data from LEP-II should show definitive evidence for the heavier bottom squark if $m_2 < 120$ GeV.
- Numbers of events in $e^+e^- \rightarrow \tilde{b}_1\tilde{b}_2^*$ vs m_2 for 3 LEP-II energies.



Polarization of J/ψ 's at the Tevatron

- INTRODUCTION:

- Gluon fragmentation into J/ψ is the dominant production mechanism at large p_T at the Tevatron.
- The gluon produces a $Q\bar{Q}$ pair that evolves nonperturbatively into the J/ψ .
- Nonrelativistic QCD (NRQCD) velocity-scaling rules predict that, in the evolution, the spin-non-flip interactions dominate over spin-flip interactions.
 - Corrections of order $v^2 \approx 0.3$.
- Hence, the J/ψ is predicted to take on most of the transverse polarization of the gluon (P. Cho, M. Wise).
- The CDF data for the polarization parameter α lie significantly below the prediction at the largest p_T :



Relativistic Corrections to Gluon Fragmentation into J/ψ

G. Bodwin and J. Lee

Phys. Rev. D **69**, 054003 (2004)

- **MOTIVATION:**

- Relativistic corrections to quarkonium decay processes are known to be large.
- Investigate whether large relativistic corrections can affect the predicted net transverse polarization of J/ψ 's at the Tevatron.


- **COMPUTATION:**

- Compute in perturbation theory the short-distance coefficients for the fragmentation process.
- Estimate the nonperturbative NRQCD matrix elements from the the quarkonium binding energy (Gremm-Kapustin relation).
- Computation of the fragmentation in the color-octet channel is simple (two-body phase space).

- Computation of the fragmentation in the color-singlet channel is more involved:
 - Three-body phase space: carry out two out of four nontrivial integrations analytically using subtle changes of variables.
 - Very large expressions manipulated using Mathematic, Reduce.

- **RESULTS:**

- Confirmed previous calculations of leading-order fragmentation short-distance coefficients (E. Braaten, T. C. Yuan, J. Lee).
- Computed fragmentation short-distance coefficients in order v^2 :



The color-singlet short-distance coefficients $d_1(z)$ and $d'_1(z)$.

The scaling factors in this figure are $c_1 = 10^{-4} \times \alpha_s^3/m^3$ and

$c'_1 = 10^{-3} \times \alpha_s^3/m^3$.

- Correction to color-singlet short-distance coefficient:
 $2.45v^2 \approx 70\%$ for J/ψ .
- Correction to color-octet short-distance coefficient:
 $-1.8v^2 \approx -50\%$ for J/ψ .
- The change in the color-singlet short-distance coefficient reduces the predicted α by about 10% at the largest p_T .
 - Still disagrees with the CDF data point at the highest p_T .
- The change in the color-octet short-distance coefficient affects the size of the color-octet matrix element in the fit to the Tevatron data.
 - No change in the predicted production rate at the Tevatron.
 - Will eventually be important in testing universality of matrix elements from process to process.

Lattice Computation of Spin Correlations in NRQCD Color-Octet Matrix Elements

G. Bodwin, J. Lee, D. Sinclair

- MOTIVATION:

- Would like to test the validity of the NRQCD velocity-scaling rules for the J/ψ production matrix elements.
- It is not known how to formulate the computation of production matrix elements on a Euclidean lattice.
- Instead, test the velocity-scaling rules on the corresponding decay matrix elements.

- PROGRESS TO DATE:

- Two versions of the required lattice code have been written independently and now agree in their numerical output.
- Modifications to the lattice action are being investigated.
- Comparisons with published results for quarkonium lattice masses from the NRQCD Collaboration are ongoing.

- PRELIMINARY RESULTS:

- Spin flips from a spin-triplet state to a spin-singlet state are large.
 - Suggests that the η_c production rate at the Tevatron may be comparable to the J/ψ production rate.
- Spin flips from a transversely polarized state to a longitudinally polarized state are small.
 - Suggests that the prediction of large transverse polarization at large p_T at the Tevatron will hold up.
- Further cross checking, production running, and analysis of the lattice data are expected to take several months.

- Stay tuned!

LATTICE QCD — D. K. Sinclair.


- Charmonium decay matrix-elements related to high- p_T charmonium production and polarization. (with G. T. Bodwin and J. Lee).
- The transition from hadronic-matter to a quark-gluon plasma at finite temperature and small densities and the search for the critical endpoint. (with J. B. Kogut).
- The transition from hadronic-matter to a quark-gluon plasma at finite temperature at zero quark mass. (with J. B. Kogut).
- The spectrum of lattice QCD at zero quark mass and tests of chiral extrapolation. (with J. B. Kogut).

Comments

- QCD finite T transition relevant to early universe, high energy heavy-ion and possibly hadron collisions. Yields information on chiral symmetry breaking, confinement and QCD dynamics in general.
- Chiral extrapolation is a persistent source of poorly controlled errors in lattice QCD calculations.
- Computing is performed on IBM SP's at NERSC and NPACI, and Linux clusters at LCRC (Argonne) and NCSA.


Lattice QCD at finite T and μ_I

- QCD at finite quark-number chemical potential μ has a complex fermion determinant.
- Ignoring the phase we have QCD at finite isospin (I_3) chemical potential $\mu_I = 2\mu$, which can be simulated.
- For $N_f = 2$, T_c decreases *slowly* with increasing μ These simulations are being extended to $N_f = 3$
- $N_f = 3$ simulations are in progress.



Lattice QCD with massless quarks at finite T

- The transition from hadronic matter to a quark-gluon plasma for 2 light quark flavours is a crossover for $m_q \neq 0$.
- It becomes a second-order chiral-symmetry-restoring transition at $m_q = 0$.
- Lattice QCD action improved to enable simulating at $m_q = 0$.
- An analysis of the behaviour of the chiral order parameter in the neighbourhood of this transition on $16^3 \times 8$ and $24^3 \times 8$ lattices ($\gamma = 3/\lambda^2$) has been performed.




Hadron spectrum in chiral limit under investigation

Beautiful Mirrors and Collider Phenomenology



D. Morrissey and C.E.M. Wagner, Phys. Rev. D 69, 053001 (2004)

- Discrepancy between value of $\sin^2 \theta_W$ obtained from lepton and bottom-quark asymmetries
- It may be arranged by the introduction of heavy vector $SU(2)$ doublet and singlet b-like quarks.
- Modification of Z -coupling due to mixing with b-quark. Masses of heavy top-like and bottom-like quarks smaller than 250 GeV and 300 GeV, respectively.
- They may be found at the Tevatron with $4 fb^{-1}$.
- Improvement of Unification conditions



Heavy Quark Impact on Higgs Physics

- Extra heavy quarks enhance the Higgs coupling to gluons, and therefore the dominant Higgs production channel at hadron colliders.
- Branching ratio of decay into tau leptons also enhanced.




Associated Production of a Z Boson and a Heavy-Quark Jet


J. Campbell, R. K. Ellis, F. Maltoni and S. Willenbrock

[hep-ph/0312024], accepted for publication in Phys. Rev. D

- **NLO calculation** using heavy-quarks in the initial state.



- Background at the Tevatron and LHC to $gb \rightarrow hb$.
- At the LHC, measures heavy-quark distribution functions.
 - no direct measurement at present
 - needed for $b\bar{b} \rightarrow h$, $gb \rightarrow hb$, $gb \rightarrow H^-t$, single top
- Including detector efficiency, there are approximately 80 $Z(\rightarrow \ell^+\ell^-)b$ events for every 100pb^{-1} at the Tevatron.
- Can reduce systematic errors by considering the ratio of tagged $Z + b$ to untagged $Z + \text{jet}$.



- **theory** prediction for ratio is 0.02 ± 0.04
- **D0** Run II measurement is 0.024 ± 0.007

Brandon Murakami

- Investigated theoretical expectations for lepton flavor violation (LFV) in rare muon decay and rare tau decay.
- Under the minimal assumptions of supergravity and neutrino oscillations, the lightest sparticle may account for the cold dark matter (CDM).
- The non-observation of sparticle CDM is consistent with the non-observation of $\mu \rightarrow e\gamma$, while simultaneously predicting probable observation at next generation $\mu \rightarrow e\gamma$ experiments.
- The observation of $\tau \rightarrow \mu\gamma$ to be unlikely in near future experiments.
- Mainstream theories that explain electroweak symmetry breaking exhibit LFV, in general.
- Only supersymmetry provides a distinct correlation $\text{BR}(\mu \rightarrow e\gamma) \propto \text{BR}(\mu - e)$ between rates of $\mu \rightarrow e\gamma$ and $\mu - e$ conversion in atomic nuclei.

θ_1 is a neutrino seesaw parameter.

$\mu \rightarrow e \gamma$

branching ratio predictions

Motivation

Theory

Experiment

Expectations

General

Technicolor

Little Higgs

Extra dimensions

Supersymmetry

Conclusions

- Case: Hierarchical neutrino mass scales.
- **WMAP compliant:** $m_0 = 100 \text{ GeV}$, $m_{1/2} = 500 \text{ GeV}$, $A_0 = 0$, $\tan\beta = 10$, $\mu > 0$
- mSUGRA predicts observable $\mu \rightarrow e \gamma$.
- $\mu \rightarrow e \gamma$ can probe neutrino parameters.

Expectations from forthcoming lepton flavor violation experiments
Brandon Murakami, ANL

$\tau \rightarrow \mu \gamma$

Motivation

Theory

Experiment

Expectations

General

Technicolor

Little Higgs

Extra dimensions

Supersymmetry

Conclusions

- Belle, BaBar, and LHC may not see tau LFV.
- Case A: Hierarchical neutrino mass scales.
- **WMAP compliant:** $m_0 = 100 \text{ GeV}$, $m_{1/2} = 500 \text{ GeV}$, $A_0 = 0$, $\tan\beta = 10$, $\mu > 0$


Expectations from forthcoming lepton flavor violation experiments
Brandon Murakami, ANL

Dark Matter and Electroweak Baryogenesis

C. Balázs, M. Carena and C.E.M. Wagner, hep-ph/0403224

- Explaining baryon asymmetry and dark matter simultaneously.
- MSSM provides new CP-violating sources and an excellent dark matter candidate.
- Computation of neutralino relic density with light stops and Higgses.

MSSM



Legend:

\times $m_{Z1} < 46$ GeV $+$ $m_{W1} < 103.5$ GeV

\times stop LSP \square $\Omega h^2 > 0.129$


\blacksquare $0.095 < \Omega h^2 < 0.129$

$\sigma_{si} = \underline{1E-06}$ $\underline{1E-07}$ $\underline{1E-08}$ pb


$m_{Z1} = \underline{160}$ $\underline{140}$ $\underline{120}$ GeV

$m_{t1} = \underline{172}$ $\underline{176}$ $\underline{181}$ GeV

- Scenario is testable at the Tevatron ...




- ... and in direct dark matter detection experiments.



Dark Matter and Electroweak Baryogenesis in the nMSSM


A. Menon, D. Morrissey and C.E.M. Wagner, hep-ph/0404184

- Minimal Extension of the MSSM, with an extra singlet superfield
- Gives a natural explanation of size of μ parameter, without inducing dangerous domain walls
- First order phase transition induced already at tree level (no need of light stops)
- Same CP-violating sources as in the MSSM
- Light Higgsino-Singlino state gives an excellent dark matter description



Higgs Physics

- Large values of charged Higgs mass
- Singlet CP-even and CP-odd component lighter than 250 GeV, due to requirement of phase transition strength



- Lightest Higgs decays mostly invisibly
- Discovery difficult at Tevatron. At the LHC, feasible in weak boson fusion channel


$$\mathcal{L}_{95\%} \simeq \frac{1.2 \text{ fb}^{-1}}{\eta^2}, \quad \mathcal{L}_{5\sigma} \simeq \frac{8.0 \text{ fb}^{-1}}{\eta^2}, \quad (1)$$

Warped GUT: Towards a realistic model


Géraldine Servant

- AdS_5 geometry generates the exponential hierarchy between M_{Pl} and M_{EW} (no need for SUSY).
- All SM fields propagate in a slice of AdS_5 except the Higgs which is localized at one boundary.
- Geometrical solution to the hierarchical flavor structure of SM fermion masses.
- Unification of gauge couplings at high scale (usual M_{GUT}) despite the infinite tower of Kaluza-Klein states.
- We addressed the pb of baryon number violation and showed that a gauged baryon number symmetry can be consistent with higher dimensional GUT.
- Like in SUSY, resolving the proton stability problem leads to a stable particle: The lightest Kaluza-Klein excitation of the right-handed neutrino in $SO(10)$, which is no more sterile but interacts with TeV mass KK excitations of $SO(10)$ gauge bosons. Thus it behaves as a typical WIMP and is an ideal dark matter (DM) candidate.
- Model will soon be tested at DM direct detection expts:


Predictions for elastic scattering expts:



We predict other light KK states with high potential for discovery at colliders:



Relic density predictions



Cosmas Zachos

- Branes are often invoked in current HEP theory, but poorly understood
- In contrast to strings, brane quantization **has never been demonstrated**
- In a significant work,
C. Zachos, Phys Lett **B570** (2003) 82-88,
“Membranes and Consistent Quantization of Nambu
Dynamics ”; and with T. Curtright, [hep-th/0312048] ,
have now achieved quantization of the special brane class
of **Topological Open Membranes**
- Cosmas has also been active investigating phenomena
relevant for phenomenology.

MEMBRANES & CONSISTENT QUANTIZATION OF NAMBU DYNAMICS

T Curtright & C Zachos

[hep-th/0312048]

BRANES, STRINGS, & ODD QUANTUM NAMBU BRACKETS

- In contrast to strings, brane quantization is an open problem.

Partial gauge fixing of covariant branes yields topological open membranes (analogous to WZW interactions):

$$S = \int (z^1 \wedge dz^2 \wedge dz^3 \wedge \dots dz^p + I_1 dt \wedge dI_2 \wedge \dots dI_{p-1}),$$

(p-2)-branes undulating in p-dim spacetimes.

Their classical eqns of motion amount to a Jacobian,

$$\frac{dz^l}{dt} = \epsilon^{lij\dots k} \partial_i I_1 \partial_j I_2 \dots \partial_k I_{p-1} = \frac{\partial(z^l, I_1, I_2, \dots, I_{p-1})}{\partial(z^1, z^2, \dots, z^p)}.$$

So, classically, instead of Hamilton's eqns, they are

$$\dot{z}^l = \{z^l, I_1, I_2, \dots, I_{p-1}\},$$

the celebrated **Nambu Bracket**, with p-1 "Hamiltonians" I_i .

A p-even Nambu Bracket is the Pfaffian of the matrix of Poisson Brackets, $\{I_i, I_j\}$, and so can **always be resolved** in terms of strings of them!

- But what are the corresponding quantum equations?

A 30 year old conundrum: **reputed inconsistencies** of Quantum Nambu Brackets.

- CZ & TC proved that **Quantization is consistent for even brackets**— in fact, Maximally Superintegrable Systems in phase space are automatically quantized this way (multi-oscillator systems, Hydrogen atom, hyperspherical/chiral models,...), given **sound understanding of their NB structure**.
 - Odd QNBs do have genuine problems, essentially because they lack the correct classical limit to odd classical NBs; so they must be redefined!
 - Even QNBs are associative, and do satisfy the proper antisymmetric **Generalized Jacobi Identity**, but not a misperceived “Fundamental” Identity, adopted by mathematicians virtually without exception. That one reflects a derivation property cleverly **bypassed by all** known solved systems through **entanglement of operators**: Nature chooses associativity over this derivation property.
- ↪ Odd NBs are quantized by **embedding them** in an even space with an extra dynamical variable, promoting them to even ones, and then quantizing those, without problems.
- Systematic success in a broad class of models, including non-hamiltonian systems.

“Noncommutativity as a Possible Origin of the Ultrahigh-Energy Cosmic Ray and the TeV Photon Paradoxes”

• Extragalactic cosmic rays with $E > 4 \times 10^{19} eV$ should be stopped by microwave background ($10^{-3} - 10^{-4} eV$) photons (GKZ): $p + \gamma \rightarrow p + \pi$; and ultra-energetic γ s with $E \sim 10 - 20 TeV$ from BL Lac blazars expected to be stopped by IR background ($10^{-1} eV$), $\gamma + \gamma \rightarrow e^+ + e^-$.

• Tiny hypothesized violations of Lorentz invariance (e.g., in noncommutative and q-deformed settings) might shift the relevant thresholds and allow paradoxical events by **modifying the energy-momentum dispersion law**, $E = \sqrt{m^2 + p^2}$.

↪ Demonstration that a class of such models (Chen & Yang) cannot work:

- The underlying theory does **not** really dictate the modified dispersion law proposed.
- The dispersion utilized is plagued by **tachyons** and so violates causality/positivity.
- Such photons would usually decay in flight by themselves, $\gamma \rightarrow e^+ + e^-$.
- The data are not that paradoxical: at the most, punch-through—but the relevant thresholds are not displaced.

Additional Topics

Work has been done in other subjects, not discussed in this talk:

J. S. Lee, A. Pilaftsis, M. Carena, S. Y. Choi, M. Drees,
J. R. Ellis and C. E. M. Wagner,

“CPsuperH: A computational tool for Higgs phenomenology in the minimal supersymmetric standard model with explicit CP violation,”

Comput. Phys. Commun. **156**, 283 (2004)

M. Carena, A. Delgado, E. Ponton, T. M. P. Tait and
C. E. M. Wagner,

“Precision electroweak data and unification of couplings in warped extra dimensions”,

Phys. Rev. D **68**, 035010 (2003)

E. W. Kolb, G. Servant and T. M. P. Tait,

“The radionactive universe,”

JCAP **0307**, 008 (2003)

Present Situation and Outlook

- Theory Group has kept a vigorous and broad research program.
- In the last year theory group has been highly productive in many relevant areas of Physics.
- Efforts are being made to keep a high profile in the HEP program, emphasizing interactions with the Argonne experimenters working at the Tevatron, LHC and in Neutrino Physics.
- Very important: Establishment of a **Junior Scientist Program**, allowing good young people to stay for a long term.